

EVERY **18** MINUTES
A BRAIN ANEURYSM
RUPTURES

EARLY INTERVENTION CHANGES OUTCOMES.

A brain aneurysm is a bulge in a weakened area of a blood vessel in or around the brain.



Dr. Abdullah Hakoun

Assistant Professor of Neurology
The University of Tennessee Health Science Center

BRAIN ANEURYSM AWARENESS

MATTERS

6 WARNING SIGNS

-  Localized headache
-  Dilated pupils
-  Blurred or double vision
-  Difficulty Speaking
-  Pain above or behind the eye
-  Weakness and numbness

**A sudden, severe headache
may be a medical emergency.
call 911**



"I had no idea what the signs were...
LaLa Speaks Foundation gave me the
knowledge to feel more prepared
while advocating for my loved one."

Britteney,
Family Advocate & Summit
Participant 2026 

DID YOU KNOW

- 1 in 50 people in the U.S. has an unruptured brain aneurysm.
- About 30,000 brain aneurysm ruptures occur in the U.S. each year.
- Women are more likely than men to have a brain aneurysm, at a 3.2 ratio.
- Women over age 55 have about 1.5 times the rupture risk of men.
- African Americans and Hispanics are about twice likely to experience a rupture as Caucasians.
- Most unruptured brain aneurysms cause no symptoms and may remain undetected.
- Brain aneurysms are most prevalent between ages 35-60, but can occur at any age.

Sources: Brain Aneurysm Foundation; Mayo Clinic; Cleveland Clinic.
Reviewed August 2026. Educational information only.

SCAN TO LEARN, SHARE & ACT

-  Share the signs
-  Invite education
-  Join the campaign



LEARN, SHARE, ACT!



Follow us @lalaspeaksfoundation



lalaspeaksfoundation.org